

Contents

VOLUME 137, NUMBER 1, SEPTEMBER 1989

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| CHAULOFF, F. Physical exercise and brain monoamines: a review | 1 |
| VØLLESTAD, N. K., BLOM, P. C. S. & GRØNNERØD, O. Resynthesis of glycogen in different muscle fibre types after prolonged exhaustive exercise in man | 15 |
| KLITGAARD, H., AUSONI, S. & DAMIANI, E. Sarcoplasmic reticulum of human skeletal muscle: age-related changes and effect of training | 23 |
| JUEL, C. & WIBRAND, F. Lactate transport in isolated mouse muscles studied with a tracer technique – kinetics, stereospecificity, pH dependency and maximal capacity | 33 |
| EKMEHAG, B. L. Electrical and mechanical responses to inhibition of cell respiration in vascular smooth muscle of the rat portal vein | 41 |
| ÅRHEM, P. & JOHANSSON, S. A model for the fast 4-aminopyridine effects on amphibian myelinated nerve fibres. A study based on voltage-clamp experiments | 53 |
| IVERFELDT, K., SERFÖZÖ, P., DIAZ ARNESTO, L. & BARTFAI, T. Differential release of coexisting neurotransmitters: frequency dependence of the efflux of substance P, thyrotropin releasing hormone and [³ H]serotonin from tissue slices of rat ventral spinal cord | 63 |
| SHYU, B. C., OLAUSSON, B., HUANG, K. H., WIDERSTRÖM, E. & ANDERSSON, S. A. Effects of sympathetic stimulation on C-fibre responses in rabbit | 73 |
| SHYU, B. C., OLAUSSON, B. & ANDERSSON, S. A. Sympathetic and noradrenaline effects on C-fibre transmission: single-unit analysis | 85 |
| PERSSON, B.-E., SAKAI, T., EKBLOM, M. & MARSH, D. J. Effect of bumetanide on tubuloglomerular feedback in <i>Necturus maculosus</i> | 93 |
| SOKRAB, T.-E. O. & JOHANSSON, B. B. Regional cerebral blood flow in acute hypertension induced by adrenaline, noradrenaline and phenylephrine in the conscious rat | 101 |
| JOHANSSON, O. & SANDBERG, G. Effect of the neuropeptides β -MSH, neurotensin, NPY, PHI, somatostatin and substance P on proliferation of lymphocytes <i>in vitro</i> | 107 |
| SAND, O., CHEN, B., LI, Q., KARLSEN, H. E., BJØRO, T. & HAUG, E. Vasoactive intestinal peptide (VIP) may reduce the removal rate of cytosolic Ca^{2+} after transient elevations in clonal rat lactotrophs | 113 |
| IVERSEN, P. O. & NICOLAYSEN, G. Heterogeneous blood flow distribution within single skeletal muscles in the rabbit: role of vasomotion, sympathetic nerve activity and effect of vasodilation | 125 |
| BATRA, S. & ANDERSSON, K.-E. Oestrogen-induced changes in muscarinic receptor density and contractile responses in the female rabbit urinary bladder | 135 |
| AUGUSTINSSON, O., JÓNASSON, H. & JUNKERGÅRD, J. Aldosterone secretion during acute metabolic and respiratory alkalosis in the goat | 143 |

Rapid communications

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| STJÄRNE, L., MSGHINA, M. & STJÄRNE, E. K^+ channel-blocking agents may augment sympathetic transmitter secretion by action 'upstream' of varicosities | 151 |
| BAO, J. X. & STJÄRNE, L. Field stimulation-induced, endothelium-dependent relaxation of rat tail artery is blocked by α,β -methylene ATP and known inhibitors of the arachidonic acid cascade | 153 |

| | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| FUXE, K., CINTRA, A., ANDBJER, B., ÄNGGÅRD, E., GOLDSTEIN, M. & AGNATI, L. F. Centrally administered endothelin-1 produces lesions in the brain of the male rat | 155 |
| FUXE, K., ANDBJER, B., KALIA, M. & AGNATI, L. F. Centrally administered endothelin-1 produces apnoea in the α -chloralose-anaesthetized male rat | 157 |

VOLUME 137, NUMBER 2, OCTOBER 1989

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| ØSTENSEN, J. Analysis of reabsorbate concentrations in the proximal tubules of dog kidneys, during osmotic diuresis | 163 |
| ØSTENSEN, J., STOKKE, E. S., BUGGE, J. F., LANGBERG, H. & KIIL, F. Difference between hypertonic NaCl and NaHCO ₃ as osmotic diuretics in dog kidneys | 177 |
| ØSTENSEN, J., STOKKE, E. S., HARTMANN, A., WENSELL, K. & KIIL, F. Low oxygen cost of carbonic anhydrase-dependent sodium reabsorption in the dog kidney | 189 |
| PARK, R. G., CLEVERS, J., MCKINLEY, M. J. & RUNDGREN, M. Renal denervation does not prevent dehydration-induced natriuresis in sheep | 199 |
| BUSTAMANTE, S. A., JODAL, M., NILSSON, N. J. & LUNDGREN, O. Evidence for a counterpart exchanger in the intestinal villi of suckling swine | 207 |
| DIENER, M. & RUMMEL, W. Actions of the Cl ⁻ channel blocker NPPB on absorptive and secretory transport processes of Na ⁺ and Cl ⁻ in rat descending colon | 215 |
| EKBLAD, E. B. M. Omeprazole: inhibitor of both acid formation and translocation in gastric mucosa | 223 |
| LARSSON, O. & OLGART, L. The enhancement of carbachol-induced salivary secretion by VIP and CGRP in rat parotid gland is mimicked by forskolin | 231 |
| OHLSOHN, L. & LINDSTRÖM, P. Growth hormone-releasing factor induces a biphasic modulation of K ⁺ permeability in perfused cultured rat somatotrophs | 237 |
| HELIG, M., VECSEI, L. & WIDERLÖV, E. Opposite effects of centrally administered neuropeptide Y (NPY) on locomotor activity of spontaneously hypertensive (SH) and normal rats | 243 |
| OLSSON, K., DAHLBORN, K., NYGREN, K., KARLBERG, B. E., ANDÉN, N.-E. & ERIKSSON, L. Fluid balance and arterial blood pressure during intracarotid infusions of atrial natriuretic peptide (ANP) in water-deprived goats | 249 |
| ANDERSSON, S. E. Effects of intravenous calcitonin gene-related peptide (CGRP) on local blood flow in the cat | 259 |
| FRANCO-CERECEDA, A. & LUNDBERG, J. M. Post-occlusive reactive hyperaemia in the heart, skeletal muscle and skin of control and capsaicin-pre-treated pigs | 271 |
| ANDERSSON, S. E. Cardiovascular responses to intravenous calcitonin gene-related peptide (CGRP) in the albino rabbit | 279 |
| DAGBJARTSSON, A., HERBERTSSON, G., STEFANSSON, T. S., KJELD, M., LAGERCRANTZ, H. & ROSEN, K. G. Beta-adrenoreceptor agonists and hypoxia in sheep fetuses | 291 |
| BAER, R. & EIKEN, O. Effects of continuous positive- and negative-pressure breathing on the pattern of breathing in man during exercise | 301 |

Rapid communications

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| VON EULER, G., MEISTER, B., HÖKFELT, T. & FUXE, K. Intraventricular injection of neurotensin reduces the binding of dopamine D-2 receptors in the rat forebrain | 309 |
| STJÄRNE, L., MSGHINA, M., STJÄRNE, E. & TANAKA, E. Temperature dependence of 'local' and 'upstream' regulation of sympathetic transmitter secretion in mouse vas deferens | 311 |
| STJÄRNE, L., MSGHINA, M. & STJÄRNE, E. Is 'upstream' control of sympathetic transmitter release dependent on the length of varicosity strings? | 313 |

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| STJÄRNE, L., MSGHINA, M. & STJÄRNE, E. Individual variations in potency of α,β -methylene ATP as inhibitor of the excitatory junction current in mouse vas deferens caused by sympathetic nerve stimulation | 315 |
| PERNOW, J., HEMSÉN, A. & LUNDBERG, J. M. Increased plasma levels of endothelin-like immunoreactivity during endotoxin administration in the pig | 317 |
| YARNITSKY, D. & OCHOA, J. L. Sensations conducted by large and small myelinated afferent fibres are lost simultaneously under compression-ischaemia block | 319 |
| SAHLGREN, B. Regulation of glomerular angiotensin II receptors in hypertensive salt-sensitive Dahl rats: absence of response to variations in salt intake | 321 |

VOLUME 137, NUMBER 3, NOVEMBER 1989

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| FAGIUS, L., KARHUVAARA, S. & SUNDLÖF, G. The cold pressor test: effects on sympathetic nerve activity in human muscle and skin nerve fascicles | 325 |
| THOMSEN, C., JENSEN, K. E., ASTRUP, A., BÜLOW, J. & HENRIKSEN, O. Changes of high-energy phosphorous compounds in skeletal muscle during glucose-induced thermogenesis in man. A ^{31}P MR spectroscopy study | 335 |
| HELLSTEN WESTING, Y., EKBLOM, B. & SJÖDIN, B. The metabolic relation between hypoxanthine and uric acid in man following maximal short-distance running | 341 |
| MORITANI, T., ODDSSON, L., THORSTENSSON, A. & ÅSTRAND, P.-O. Neural and biochemical differences between men and young boys during a variety of motor tasks | 347 |
| FÄNDRIS, L., JÖNSEN, C. & LISANDER, B. Hypothalamic inhibition of duodenal alkaline secretion via a sympatho-adrenergic mechanism in the rat | 357 |
| HEMLIN, M. Interaction between sympathetic nerve fibres and epithelial transport in the rat jejunal mucosa <i>in vivo</i> | 365 |
| DAHLSTRAND, C., THEODORSSON, E., DAHLSTRÖM, A. & AHLMAN, H. VIP antisera inhibit the relaxatory motor responses of the feline sphincter of Oddi and gall-bladder induced by VIP or vagal nerve stimulation | 375 |
| KJÆLDGAARD, A., KJÆLDGAARD, M. & GAFFNEY, P. Presence of a fast-acting specific inhibitor of plasminogen activator in human parotid saliva | 379 |
| THOLLANDER, M., HELLSTRÖM, P. M. & SVENSSON, T. H. Suppression of small intestinal motility and morphine withdrawal diarrhoea by clonidine: peripheral site of action | 385 |
| ENGSTRÖM, K. G. & SANDSTRÖM, P.-E. Volume regulation in mouse pancreatic islet cells as studied by a new technique of microperfusion | 393 |
| WIKLUND, N. P., ÖHLÉN, A., WIKLUND, C. U., CEDERQVIST, B., HEDQVIST, P. & GUSTAFSSON, L. E. Neuromuscular actions of endothelin on smooth, cardiac and skeletal muscle from guinea-pig, rat and rabbit | 399 |
| RYDQVIST, B. & ZHOU, J.-Y. Potential-dependent potassium currents in the slowly adapting stretch receptor neuron of the crayfish | 409 |
| PERNOW, J. Characterization of the cardiovascular actions of endothelin <i>in vivo</i> : comparisons with neuropeptide Y and angiotensin II | 421 |
| JONASSON, H., HJELMQVIST, H. & RUNDGREN, M. Repeated hypotension induced by nitroprusside and haemorrhage in sheep: effects on vasopressin release and recovery of arterial blood pressure | 427 |
| SJÖBLOM-WIDFELDT, N., NILSSON, H., GUSTAFSSON, H., ELY, D. L. & FOLKOW, B. Effects of pre-junctional α -receptor blockade on vascular neuro-effector characteristics in rats on high sodium intake | 437 |

| | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| SJÖBLOM-WIDFELDT, N. & NILSSON, H. Calcium dependency of the post-stimulatory potentiation of the neurogenic response in small mesenteric arteries from the rat | 443 |
| APPELGREN, L.-E., ODLIND, B. & WISTRAND, P. J. Tissue distribution of ^{125}I -labelled carbonic anhydrase isozymes I, II and III in the rat | 449 |

Rapid communications

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| FRANCO-CERECEDA, A., LOU, Y.-P. & LUNDBERG, J. M. Ruthenium red differentiates between capsaicin and nicotine effects on cardiac sensory nerves | 457 |
| ILLES, P. & REGENOLD, J. T. ω -Conotoxin GVIA and nifedipine inhibit Ca^{2+} action potentials in rat locus coeruleus neurons | 459 |
| MEISTER, B., VILLAR, M. J., SCHALLING, N., EHRLICH, M., GREENGARD, P. & HÖKFELT, T. Demonstration of DARPP-32 in pituitary cells of the neurohypophysis – decreased expression after administration of hyperosmotic stimuli | 461 |
| XU, X.-J., WIESENFELD-HALLIN, Z., VILLAR, M. J. & HÖKFELT, T. Intrathecal galanin antagonizes the facilitatory effect of substance P on the nociceptive flexor reflex in the rat | 463 |
| PELTO-HUIKKO, M., PERSSON, H., SCHALLING, N. & HÖKFELT, T. Immunocytochemical demonstration of cholecystokinin-like immunoreactivity in monkey testis and epididymis | 465 |
| SCHALLING, M., FRIBERG, K., BIRD, E., GOLDSTEIN, M., SCHIFFMANN, S., MALLIEUX, P., VAN DER HAEGHEN, J.-J. & HÖKFELT, T. Presence of cholecystokinin mRNA in dopamine cells in the ventral mesencephalon of a human with schizophrenia | 467 |

VOLUME 137, NUMBER 4, DECEMBER 1989

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| BRIDÉN, S. Electrophysiological characteristics of the <i>Necturus</i> proximal duodenal mucosa: effects of ion substitutions | 469 |
| BRIDÉN, S. & FLEMSTRÖM, G. Alkaline secretion by <i>Necturus</i> proximal duodenal mucosa | 481 |
| ALBERTS, P. Effects of $\text{N}^6,2'$ - <i>O</i> -dibutyryladenosine 3',5'-cyclic monophosphate, adenosine, and of oxotremorine and 3-isobutyl-1-methylxanthine on the electrically evoked [^3H]acetylcholine secretion in the guinea-pig myenteric plexus | 489 |
| MOLGÓ, J., DASGUPTA, B. R. & THESLEFF, S. Characterization of the actions of botulinum neurotoxin type E at the rat neuromuscular junction | 497 |
| SHYU, B. C., OLAUSSON, B. & RYDENHAG, B. Field potential analysis of the cortical projection of the central lateral nucleus in the cat | 503 |
| LUNDVALL, J. & LÄNNE, T. Large capacity in man for effective plasma volume control in hypovolaemia via fluid transfer from tissue to blood | 513 |
| KARLSTRÖM, G., ARNAM, V., BERGSTRÖM, G., MUIRHEAD, E. E., RUDENSTAM, J. & GÖTHBERG, G. Renal and circulatory effects of medullipin I, as studied in the <i>in-vivo</i> cross-circulated isolated kidney and intact Wistar-Kyoto (WKY) rat | 521 |
| SONG, Y.-H. & MÅRDH, S. The occurrence of gastric and duodenal auto-antibodies in peptic ulcer disease | 535 |

Rapid communications

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| HAEGERSTRAND, A., HEMSEN, A., GILLIS, C., LARSSON, O. & LUNDBERG, J. M. Endothelin: presence in human umbilical vessels, high levels in fetal blood and potent constrictor effect | 541 |
| NIE, Z. T. & HENRIKSSON, J. <i>In-vitro</i> stimulation of the rat epitrochlearis muscle. III. Endogenous levels of branched-chain amino acids are maintained during acute contractions even in the absence of an exogenous supply | 543 |

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| OHLSSON, L., LINDSTRÖM, P. & SEHLIN, J. Perchlorate stimulates potassium-induced growth hormone release in cultured rat somatotrophs | 545 |
| VAN DEN BOS, R., COOLS, A. R. & ÖGREN, S. O. Neurokinin A enhances the stimulatory effects of d-amphetamine on motor activity in the nucleus accumbens of the rat | 547 |
| CANLON, B., CARTAUD, J. & CHANGEUX, J. P. Localization of α -bungarotoxin binding sites on outer hair cells from the guinea-pig cochlea | 549 |
| FUXE, K., TINNER, B., JANSON, A. M., CINTRA, A., STAINES, W. & AGNATI, L. F. On the cellular localization and distribution of the ganglioside GM1 in the rat brain as revealed by immunofluorescence histochemistry of cholera toxin binding sites | 551 |
| PERNOW, J. & LUNDBERG, J. M. Endothelin-like immunoreactivity is released from the pig spleen during asphyxia | 553 |
| HÖKFELT, T., POST, C., FREEDMAN, J., LUNDBERG, J. M. & TERENIUS, L. Endothelin induces spinal lesions after intrathecal administration | 555 |
| CINTRA, A., FUXE, K., ÄNGGÅRD, E., TINNER, B., STAINES, W. & AGNATI, L. F. Increased endothelin-like immunoreactivity in ibotenic acid-lesioned hippocampal formation of the rat brain | 557 |
| CECCATELLI, S., DAGERLIND, Å., SCHALLING, M., WIKSTRÖM, A.-C., OKRET, S., GUSTAFSSON, J. Å., GOLDSTEIN, M. & HÖKFELT, T. The glucocorticoid receptor in the adrenal gland is localized in the cytoplasm of adrenaline cells | 559 |
| HULTING, A.-L., MEISTER, B., GRIMELIUS, L., WERSALL, J., ÄNGGÅRD, A. & HÖKFELT, T. Production of a galanin-like peptide by a human pituitary adenoma: immunohistochemical evidence | 561 |
| FUXE, K., ÄNGGÅRD, E., LUNDGREN, K., CINTRA, A., AGNATI, L. F., GALTON, S. & VANE, J. Localization of [125 I]endothelin-1 and [125 I]endothelin-3 binding sites in rat brain | 563 |
| LARSSON, P. T., OLSSON, G., HJEMDAHL, P. & NILSSON, J. Mental stress and adrenaline increase plasma growth factor activity in humans | 565 |
| ECKBERG, D. L., WALLIN, B. G., FAGIUS, J., LUNDBERG, L. & TOREBJÖRK, H. E. Prospective study of symptoms after human microneurography | 567 |
| HEDQVIST, P., RAUD, J., PALMERTZ, U., HAEGGSTRÖM, J., NICOLAOU, K. C. & DAHLÉN, S.-E. Lipoxin A ₄ inhibits leukotriene B ₄ -induced inflammation in the hamster cheek pouch | 571 |
| AUTHOR AND SUBJECT INDEX | 573 |